

GATE - 1989

www.gdc4gpat.com

PY- PHARMACEUTICAL SCIENCES

Time: 3 hours Maximum Marks: 200

- N. B. 1. This question paper contains two parts A and B.
 - 2. Answer all the question from part A.
 - 3. Answer Any 20 Question from part B.

PART - A

- N. B. 1. There are 2 sections in this part
 - 2. Answer all the question in both sections 1 and 2.
 - 3. Answer should be given serial order in the answer book.
 - 4. Do not skip question while writing the answers.
 - 5. Write the question number and show your answer by writing the alphabet (against the No.) in Capital letters.
 - 6. In section 1 each question carriers 1-Marks.
 - 7. In section 2 each question carries 2-marks.
 - 8. A model is shown at the beginning of each section in part A.
 - 9. Answer to the question in this part must be Witten in the first three pages only.

SECTION - I

CHOOSE THE CORRECT ANSWER

Model Ouestion

- 1. Repeated administration of Tyramine results in its decreasing effectiveness:
 - (a) Gets detoxicated easily
 - (b) Displaces nor-adrenaline from nerve ending binding site
 - (c) Displaces adrenaline from nerve ending binding site
 - (d) None of the above
- 2. Atropine on hydrolysis with Barium hydroxide gives:
 - (a) Tropanol and Tropic acid

(b) Scopine and Tropic acid

(c) Ecgonine and Benzoic acid

(d) Benzyl Ecgonine and Methanol

Visit - www.gdconlinetest.in | Attempt 1 Free Demo Test

Email: gdcgpat037@gmail.com

PREVIOUS
YEAR PAPER

Join All India Classroom/Online Test Series for GPAT Conducted by GPAT Discussion Center all over India. Question papers are designed by the expert faculties. It may helps you in focusing your Exampreparation.



GATE - 1989

www.gdc4gpat.com

Vi	sit - www.gdconlinetest.in Attempt 1 Free Demo Test	Email: gdcgpat037@gmail.com	ПС
	(c) Salmonella typii	(d) Escherichia coil	
	(a) Bacillus subtilis	(b) Micrococcus luteus	
12.	In the official bioassay of Erythromycin strai	in used is:	
	(c) Reserpine	(d) Polypeptides	
	(a) Digitoxin	(b) Lysergic acid derivatives	
11.	Claviceps purpurea yields after infecting ova	ries of Graminaceous plants:	
	(a) 6.5 (b) 13.8	(c) 25.0 (d) 40.0	
10.	The HLB value of sodium lauryl sulphate is:		
	(d) α - Carboxy benzyl penicillin		
	(c) β - [Hydroxy analogue of Benzyl penicill	lin	
	(b) 4 - [D-(-) α – amino p-hydroxyacetamido	o] penicillanic acid	
	(a) 6 - [D-(-) α – amino p-hydroxyacetamido	o] penicillanic acid	
9.	Chemical name of amoxicillin is:		
	(c) Precipitating steroids from solution	(d) Precipitating Anthraquinone glycosides	
	(a) Cardiac action	(b) Hypotensive action	
8.	C3 O-glycoside digitoxin is used for:	CENTER	
	(c) Vacuum drying	OISCU(d) Fluid bed drying	
	(a) Spray drying	(b) Freeze drying	
7.	For drying blood plasma the following techin	TPAI	
	(c) Digitalis and Senna glycosides	(d) Digitalis and Amygdalin	
	(a) Digitalis and squill glycosides	(b) Digitalis and strophantus glycosides	
6.	C 17 α - β unsaturated lactone ring is a comm		
	(d) An alternation in enzyme that utilize PA		
	(c) An increasing product of drug antagonis		
	(b) An alternative metabolic pathway for sy		
	(a) An increasing capacity to inactivate or d	, ,	
5.	Most accepted mechanism for developing ba	-	
	(c) Gentianaceae	(d) Polygonaceae	
	(a) Apocynaceae	(b) Scrophularaceae	
4.	Stratified cork and forked are the characteri		
	(c) 66.70% w/w	(d) 40.74% w/w	
	(a) 85% w/w	(b) 60.70% w/w	
3.	The concentration of sucrose in simple Syrup	p BP is:	

Email: gdcgpat037@gmail.com

PREVIOUS

Discussion Center all over India. Question



GATE - 1989

www.gdc4gpat.com

13.	13. The disintegration time for sugar coated tablet is						
	(a) 30 minutes (b) 45 minutes	(c) 60 minutes (d) 75 minutes					
14.	Idioblasts of crystal layer of calcium oxalate is a diagram	nostic feature of					
	(a) Hyoscyamus Niger leaves	(b) Deadly nightshade leaves					
	(c) Cinchona bark	(d) Senna leaves					
15.	Antibiotic which interacts with calcium ion is:						
	(a) Erythromycin (b) Streptomycin	(c) Tetracycline (d) Ampicillin					
16.	Flow rate of granules from the hopper can be impro	ved by adding;					
	(a) Disintegrant (b) Glidant	(c) Binder (d) Lubricant					
17.	Silicon carbide rod heated to a high temperature is u	ised as a;					
	(a) Detector in infra red spectroscope	(b) Source of light in infra re spectroscope					
	(c) Source of light fluorimetery	(d) Detector in gas chromatography					
18.	Anomocytic type stomata are found in the leaves of:						
	(a) Fox glove	(b) Urginea maritime					
	(c) Cassia acutifolia	(d) Atropa belladonna					
19.	Liver microsomal enzymes are stimulated (enzymic	induction) by:					
	(a) Cimetidine (b) Phenobarbitone	(c) Procaine (d) Adrenaline					
20.	Enteric coating is achieved by using: C E 1	NTER					
	(a) Hydroxy propyl methyl cellulose	(b) Carboxy methyl cellulose					
	(c) Cellulose acetate Phthalate	(d) Povidone					
21.	Car price reaction is applied for the photometric eva	lluation of:					
	(a) Vitamin A	(b) Tocopherol					
	(c) Nandrolone Phenyl Propionate	(d) Benzodiazepine					
22.	Peroxide enzyme present in acacia is identified by:						
	(a) Borntragers test	(b) Molisch's test					
	(c) Oxidation and extraction in Benzene	(d) Oxidation and treatment with Benzididine					
23.	Prostaglandins are a group of related:						
	(a) Alcohols (b) Aldehydes	(c) Fatty acid (d) Alkaloids					
24.	Licence to sell drug specified in schedule C and C1 is	given from number					
	(a) -19 (b) -18	(c) -21 (d) -24					
25.	Liqiude paraffin exhibits:						
	(a) Plastic flow	(b) Newtonian flow					
	(c) Pseudoplastic flow	(d) Dilatant flow					
17.	isit - www.gdconlinetest in Attempt 1 Free Demo Test	Fmail: adcapat037@amail.com					



www.gdc4gpat.com



(a) Inhibits the ovulation (b) Inhibits the implantation of the fertilized ovum (c) Inhibits the fertilization of ovum (d) Inhibits development of endometrium More of earthy matter in a Rhizome is determine by: (a) Total ash value (b) The earthy material is separated and then weighed (c) The Rhizome is washed in water and the in hydrochloric acid finally it is weighed (d) Acid insoluble ash value 8. Lidocaine is synthesized from: (a) 2:6-dimethyl-5-amino methyl benzene (c) 2:6-dimethyl-5-amino methyl benzene (d) 2-methyl-6-ethyl-5-amino methyl benzene (e) 2:6-dimethyl-5-amino methyl benzene (g) 2:6-dimethyl-5-amino methyl benzene (h) 2-methyl-6-ethyl-5-amino methyl benzene (g) 2:6-dimethyl-5-amino methyl benzene (h) 2-methyl-6-ethyl-5-amino methyl benzene (26.	Estrogenic and Progestrogenic combination mainly:								
(a) Total ash value (b) The earthy material is separated and then weighed (c) The Rhizome is washed in water and the in hydrochloric acid finally it is weighed (d) Acid insoluble ash value 28. Lidocaine is synthesized from: (a) 2:6-dimethyl-5-amino methyl benzene (b) 2:6-dimethyl-5-nitro methyl benzene (c) 2:6-xylidene (d) 2-methyl-6-ethyl-5-nitro methyl benzene (e) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene (e) 2:6-xylidene (f) 147°C (g) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragatus gummifer is depend on: (a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glucose (d) 2:6-deoxy glactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(a) Inhibits the ovulation	(b)	Inhibits the implantat	tion of the fertilized ovum					
(a) Total ash value (b) The earthy material is separated and then weighed (c) The Rhizome is washed in water and the in hydrochloric acid finally it is weighed (d) Acid insoluble ash value 28. Lidocaine is synthesized from: (a) 2:6-dimethyl-5-amino methyl benzene (c) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene (e) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene 29. Sterilization temperature for aqueous solution in autockave (Moist heat) is: (a) 72°C (b) 121°C (c) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Clofazimine + Streptomycine 31. The gummy nature Astragalus gummifer is depend on: (a) More of Methoxly group of Bassorin (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaklehyde and 2-amino hydantoin (b) 5-Nitro 2-furaklehyde and hydantoin (c) 5-amino 2-furaklehyde and 2-amino hydantoin (d) 5-Nitro 2-furaklehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10° mole/lit (b) 10° mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(c) Inhibits the fertilization of ovum	(d)	Inhibits development	of endometrium					
(b) The earthy material is separated and then weighed (c) The Rhizome is washed in water and the in hydrochloric acid finally it is weighed (d) Acid insoluble ash value 28. Lidocaine is synthesized from: (a) 2:6-diwethyl-5-amino methyl benzene (c) 2:6-dylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene (e) 2:6-dylidene (f) 121°C (g) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Clofazimine + Rifampin (e) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragalus gummifer is depend on: (a) More of hydroxyl group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10° mole/lit (b) 10° mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitalis purpurea is: (a) 2:6-deoxy glacose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (d) Propranalol	27.	More of earthy matter in a Rhizome is determine by	:							
(c) The Rhizome is washed in water and the in hydrochloric acid finally it is weighed (d) Acid insoluble ash value 28. Lidocaine is synthesized from: (a) 2:6-dimethyl-5-amino methyl benzene (c) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene (e) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene 29. Sterilization temperature for aqueous solution in autoclave (Moist heat) is: (a) 72°C (b) 121°C (c) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragalus gummlfer is depend on: (a) More of Methoxly group of Bassorin (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and hydantoin (e) 5-amino 2-furaldehyde and 2-amino hydantoin (f) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitals purpurea is: (a) 2:6-deoxy glactose (b) 2:6-deoxy glactose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glactose (d) 2:6-deoxy glactose (e) 2:6-deoxy Rhamnose (d) 2:6-deoxy glactose (d) 2:6-deoxy glactose (e) 2:6-deoxy inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(a) Total ash value								
(d) Acid insoluble ash value 28. Lidocaine is synthesized from: (a) 2:6-dimethyl-5-amino methyl benzene (b) 2:6-dimethyl-5-amino methyl benzene (c) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene 29. Sterilization temperature for aqueous solution in autoclave (Moist heat) is: (a) 72°C (b) 121°C (c) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Terracycline + Streptomycine 31. The gummy nature Astragalus gummlfer is depend on: (a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (c) 5-mino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitalis purpurea is: (a) 2:6-deoxy glacose (b) 2:6-deoxy glacose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(b) The earthy material is separated and then weig	hed							
28. Lidocaine is synthesized from: (a) 2:6-dimethyl-5-amino methyl benzene (b) 2:6-dimethyl-5-nitro methyl benzene (c) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene (e) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene (e) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene (e) 2:6-xylidene (d) 160 °C (e) 147 °C (d) 160 °C (d) 160 °C (e) 147 °C (d) 160 °C (d) 160 °C (e) 147 °C (d) 160 °C (e) 160 °C (f) 160 °C (g) 160 °C (h)		(c) The Rhizome is washed in water and the in hydrony	roch	loric acid finally it is v	veighed					
(a) 2:6-dimethyl-5-amino methyl benzene (b) 2:6-dimethyl-5-nitro methyl benzene (c) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene 29. Sterilization temperature for aqueous solution in autoclave (Moist heat) is: (a) 72°C (b) 121°C (c) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (b) Dapsone + Clofazimine + Rifampin (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragatus gummifer is depend on: (a) More of Methoxly group of Bassorin (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaklehyde and 2-amino hydantoin (b) 5-Nitro 2-furaklehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10 ⁴ mole/lit (b) 10 ⁷ mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy glacose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glacose (e) 2:6-deoxy Rhamnose (d) 2:6-deoxy glacose (d) 2:6-deoxy glacose (e) 2:6-deoxy Rhamnose (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(d) Acid insoluble ash value								
(c) 2:6-xylidene (d) 2-methyl-6-ethyl-5-amino methyl benzene 29. Sterilization temperature for aqueous solution in autoclave (Moist heat) is: (a) 72°C (b) 121°C (c) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (b) Dapsone + Clofazimine + Rifampin (d) Dapsone + Erythromycine + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragalus gumm fer is depend on: (a) More of Methoxly group of Bassorin (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaklehyde and 2-amino hydantoin (b) 5-Nitro 2-furaklehyde and hydantoin (c) 5-amino 2-furaklehyde and 2-amino hydantoin (d) 5-Nitro 2-furaklehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	28.	Lidocaine is synthesized from:								
29. Sterilization temperature for aqueous solution in autoclave (Moist heat) is: (a) 72°C (b) 121°C (c) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (b) Dapsone + Clofazimine + Rifampin (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragalus gummifer is depend on: (a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaklehyde and 2-amino hydantoin (d) 5-Nitro 2-furaklehyde and hydantoin (c) 5-amino 2-furaklehyde and 2-amino hydantoin (d) 5-Nitro 2-furaklehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be about (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glucose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(a) 2:6-dimethyl-5-amino methyl benzene	(b)	2 : 6-dimethyl-5-nitro	methyl benzene					
(a) 72°C (b) 121°C (c) 147°C (d) 160°C 30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (b) Dapsone + Clofazimine + Rifampin (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragalus gummifer is depend on: (a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaklehyde and 2-amino hydantoin (b) 5-Nitro 2-furaklehyde and hydantoin (c) 5-amino 2-furaklehyde and 2-amino hydantoin (d) 5-Nitro 2-furaklehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy Rhamnose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glactose 36. Additional of sodium chloride to sodium Oleate emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(c) 2:6-xylidene	(d)	2-methyl-6-ethyl-5-ar	nino methyl benzene					
30. Following combination is suggested in the treatment of Leprosy: (a) Dapsone + Ampicinllin + Clofazimine (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragalus gummifer is depend on: (a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	29.	Sterilization temperature for aqueous solution in au	tocla	ve (Moist heat) is:						
(a) Dapsone + Ampicinllin + Clofazimine (b) Dapsone + Clofazimine + Rifampin (c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragalus gummifer is depend on: (a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy glactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(a) 72°C (b) 121°C	(c)	147 °C	(d) 160°C					
(c) Dapsone + Erythromycin + Rifampin (d) Dapsone + Teracycline + Streptomycine 31. The gummy nature Astragalus gummifer is depend on: (a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaklehyde and 2-amino hydantoin (b) 5-Nitro 2-furaklehyde and hydantoin (c) 5-amino 2-furaklehyde and 2-amino hydantoin (d) 5-Nitro 2-furaklehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	30.	Following combination is suggested in the treatment	t of l	Leprosy:						
31. The gummy nature Astragalus gummifer is depend on: (a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-dedoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(a) Dapsone + Ampicinllin + Clofazimine	(b)	Dapsone + Clofazimii	ne + Rifampin					
(a) More of Methoxly group of Bassorin (b) The carbohydrate content (c) More of hydroxyl groups of the sugar moiety 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be about (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(c) Dapsone + Erythromycin + Rifampin	(d)	Dapsone + Teracyclir	ne + Streptomycine					
(c) More of hydroxyl groups of the sugar moiety (d) More of protein contain of the drug 32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	31.	The gummy nature Astragalus gummifer is depend o	n:	SION						
32. The vitamin administered with isoniazid to minimize its adverse reaction is (a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be about (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(a) More of Methoxly group of Bassorin	1	I' H' R						
(a) Vitamin A (b) Pyridoxine (c) Biotin (d) Pantothenic acid 33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(c) More of hydroxyl groups of the sugar moiety	(d)	More of protein cont	ain of the drug					
33. For the synthesis of Nitrofurantoin which one of the following combination of chemicals are used: (a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be about (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-dedoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	32.	The vitamin administered with isoniazid to minimize	its	adverse reaction is						
(a) 5-Nitro 2-furaldehyde and 2-amino hydantoin (b) 5-Nitro 2-furaldehyde and hydantoin (c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be abou (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-dedoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(a) Vitamin A (b) Pyridoxine	(c)	Biotin	(d) Pantothenic acid					
(c) 5-amino 2-furaldehyde and 2-amino hydantoin (d) 5-Nitro 2-furaldehyde and barbituric acid 34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be about (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-dedoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	33.	For the synthesis of Nitrofurantoin which one of the	foll	owing combination of	chemicals are used:					
34. To get the optimum optical density of the solution for 1 cm thick layer the concentration should be about (a) 10-4 mole/lit (b) 10-7 mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		(a) 5-Nitro 2-furaldehyde and 2-amino hydantoin	(b)	5-Nitro 2-furaldehyde	and hydantoin					
(a) 10 ⁻⁴ mole/lit (b) 10 ⁻⁷ mole/lit (c) 0.1 gm/lit (d) 0.5 gm/lit 35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol										
35. The sugar moiety of Digitails purpurea is: (a) 2:6-deoxy allose (b) 2:6-dedoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	34.			•						
 (a) 2:6-deoxy allose (b) 2:6-deoxy glucose (c) 2:6-deoxy Rhamnose (d) 2:6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol 			(c)	0.1 gm/lit	(d) 0.5 gm/lit					
(c) 2 : 6-deoxy Rhamnose (d) 2 : 6-deoxy galactose 36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	35.									
36. Additional of sodium chloride to sodium Oleate emulsion will: (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol		•	. ,	, ,						
 (a) Stabilize emulsion (b) Destabilize emulsion (c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol 										
(c) Decrease the globule size of the emulsion (d) None of the above 37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol	36.									
37. Anti hypertensive drug inhibits the rennin angiotensin system is: (a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol			. ,							
(a) Reserpine (b) Captopril (c) Methyl dopa (d) Propranalol			. ,							
	37.									
		.,								

papers are designed by the expert faculties. It may helps you in focusing your Exam preparation.



www.gdc4gpat.com

GATE - 1989

38. Acidity of Ascorbic acid is due to the presence of:

(a) Free carboxylic acid

(b) A number of hydroxyl group

(c) Enolic groups

(d) None of the above

39. Progesterone injection BP is a sterile solution in:

- (a) Water
- (b) Ethyl oleate
- (c) Propylene glycol
- (d) Glycerol

40. Thiamine on treatment with sodium sulfite solution and sulfur dioxide yields:

- (a) Pyrimidine and a thiazole derivative
- (b) Pyridine and thiazole derivative
- (c) 2:3:4-Thihydropyridine and Thiophene derivatives
- (d) Pyrimidine and Thiophene derivatives.



MATCH THE FOLLOWING

2.1. Identify the correct skeleton ring present in the following compounds from the ring system listed from A to E.

- 1. Riboflavin
- 2. Estrone
- 3. Indomethacin
- 4. Nalidixic acid
- (a) 1-E, 2-A, 3-C, 4-D
- (c) 1-B, 2-C, 3-A, 4-D

- (A) Perhydro cyclopentanophenanthrene
- (B) 1:8 Naphthyridine
 - (C) Indole
 - (D) Quinolin
 - (E) Iso alloxagine
 - (b) 1-D, 2-C, 3-B, 4-A
- (d) 1-D, 2-A, 3-C, 4-B

2.2. Chosse the instrument or apparatus listed from A to E study the following:

- 1. Rheology of semi solids
- 2. Hardness of tablets
- 4. Homogenization of emulsion
- 3. Particle size in suspension
- (a) 1-D, 2-B, 3-C, 4-A
- (c) 1-D, 2-C, 3-A, 4-B

- (A) Andreasen Pipette
- (B) Monasanto tester
- (C) Ultrasonifier
- (D) Viscometer
- (E) Zeta meter
- (b) 1-E, 2-B, 3-A, 4-C
- (d) 1-C, 2-B, 3-D, 4-A



GATE - 1989

www.gdc4gpat.com

2.3.	Given below are some	microscopical	diagnostic of	the drug listed in	n A to E. Chosse	the appropriate one
------	----------------------	---------------	---------------	--------------------	------------------	---------------------

- Unlignified septate fiber
- 2. Raphides of calcium oxalate embedded in mucilage
- Anisocytic type of stomata 3.
- Star spots
- (a) 1-A, 2-B, 3-C, 4-D
- (c) 1-B, 2-C, 3-A, 4-D
- 2.4. Chosse the most appropriate drug for the following
 - 1. Potassium-sparing diuretic
 - 2. Loop diuretic
 - Osmotic diuretic
 - 4. Carbonic anhydrase inhibitor
 - (a) 1-A, 2-C, 3-E, 4-D
 - (c) 1-A, 2-C, 3-D, 4-B

- (A) Rhubarb
- (B) Solanaceous plant
- (C) Ginger
- (D) Squill
- (E) Solanaceous plants
- (b) 1-D, 2-C, 3-B, 4-A
- (d) 1-A, 2-D, 3-E, 4-A
- (A) Spiranolactone
- (B) Mannitol
- (C) Furosemide
- (D) Acetazolamide
- (E) Aldosterone
- (b) 1-A, 2-B, 3-C, 4-D
- (d) -1-C, 2-B, 3-D, 4-A
- 2.5. Transmitted colour corresponds to various wave length ranges as listed under A to E. Choose the correct wave length for the colour. CENTER
 - 1. Green
 - 2. Orange
 - 3. Yellow
 - 4. Red
 - (a) 1-A, 2-B, 3-C, 4-D
 - (c) 1-B, 2-D, 3-C, 4-E

- (A) 635-700
- (B) 520-560
- (C) 560-590
- (D) 590-635
- (E) 650-780
- (b) 1-B, 2-C, 3-A, 4-D
- (d) 1-B, 2-C, 3-D, 4-A
- 2.6. Given below equipment used in the manufacture of the following product A to E. Match them correctly.
 - Zanasi
 - 2. HEPA Filter
 - 3. Chilsonator
 - 4. Accela cota
 - (a) 1-D, 2-A, 3-C, 4-B
 - (c) 1-B, 2-C, 3-A, 4-D

- (A) Tablet granules
- (B) Tablet coating
- (C) Emulsion
- (D) Injectable
- (E) Capsules
- (b) 1-E, 2-D, 3-A, 4-B
- (d) 1-C, 2-B, 3-D, 4-A

Visit - www.gdconlinetest.in | Attempt 1 Free Demo Test

Email: gdcgpat037@gmail.com

Previous Year Paper

ALLAHABAD BILASPUR BHILAI RAIPUR NAGPUR HYDERABAD JAIPUR

papers are designed by the expert faculties. It may helps you in focusing your Exam preparation.



GATE - 1989

www.gdc4gpat.com

2.7. Match the following with the schedules listed in A to E correctly.

Requirements of factory premises

(A) P

2. Standards for disinfectant fluids (B) V

Life period of drugs

- (C) N
- 4. List of minimum equipment for the efficient running of Pharmacy
- (D) 0(E) M

(a) 1-E, 2-D, 3-A, 4-C

(b) 1-B, 2-C, 3-D, 4-A

(c) 1-B, 2-C, 3-A, 4-D

(d) 1-C, 2-B, 3-D, 4-A

2.8. Following are the reaction/tests observed in case of drugs listed in A to E. Match them correctly.

- 1. When fixed oil is exposed to U.V. rays, blue
- (A) Digoxin
- 2. On oxaidation with KMnO₄, Benzaldehyde
- (B) Benzoin
- - With ammoniacal Quaxom characteristic ballooned (C) Cinchona
- 4. Bark powder exhibits fluorescence with sulphuric acid
- (D) Palmolein

(a) 1-A, 2-B, 3-C, 4-D

(E) Gossypium barbadance

(b) 1-D, 2-B, 3-E, 4-C

(c) 1-B, 2-C, 3-A, 4-D

(d) 1-C, 2-B, 3-D, 4-A

2.9. Mechanism of Antitubercular action of the drug listed are indicate are in A to E. Choose the most appropriate DISCUSSION one.

Ethambutol

(A) Prevents the synthesis of protein and DNA and reduces R.N.A. synthesis.

2. P. A. S.

(B) Interferes with several of protein synthesis

3. Cycloserine

(C) Competitive inhibiton

4. Ethionamide

(D) Inhibits peptide synthesis in Mycobacteria

(a) 1-A, 2-B, 3-C, 4-D

(E) Inhibits DNA directed RNA Synthesis (b) 1-D, 2-A, 3-C, 4-B

(c) 1-B, 2-C, 3-A, 4-D

(d) 1-D, 2-C, 3-B, 4-A

2.10. Given below are the receptor and their antagonist (A to E). Match them correctly.

1. Histamine H₂ Receptor

(A) Atropine

2. Muscarinic Receptor

(B) Ranitidine

3. Adrenaline α receptor

(C) Pentolamine

4. Adrenaline α recptor

(D) Metaraminol

(a) 1-B, 2-A, 3-C, 4-E

(E) Metoprolol

(c) 1-B, 2-C, 3-A, 4-D

- (b) 1-D, 2-C, 3-B, 4-A
- (d) 1-C, 2-B, 3-D, 4-A

papers are designed by the expert faculties. It may helps you in focusing your Exam preparation.





www.gdc4gpat.com

2.11. Match the following regions in GIT with the pH levels indicated from A to E.

Mouth

2. Stomach

3. Deodenum

4. Large intestine

(a) 1-A, 2-D, 3-B, 4-C

(c) 1-B, 2-E, 3-D, 4-C

(A) = 5.0 - 6.0

(B) = 6.8 - 7.5

(C) = 6.8 - 7.0

(D) = 3.0 - 5.0

(E) = 1.5 - 3.0

(b) 1-A, 2-D, 3-B, 4-A

(D) 1-C, 2-B, 3-D, 4-A

2.12. Listed in A to E are some of the analytical constants. Match them correctly with the drugs given below.

1. A Leafy drug

2. A Bark

3. Eucalyptus oil

4. A fixed oil having more of unsaturated (D) Iodine value fatty acid glycerides

(a) 1-A, 2-B, 3-C, 4-D

(c) 1-E, 2-C, 3-B, 4-D

(A) Total ash value

(B) Cineole content

(C) Fibre length

(E) Stomatal index

(b) 1-D, 2-C, 3-B, 4-A

(d) 1-C, 2-B, 3-D, 4-A

2.13. Match the ingredients listed A to E with the purpose for which they are used in the formulations.

1. Film coating

2. Syrups

3. Emulsification

4. Enteric coating

(a) 1-B, 2-D, 3-A, 4-C

(c) 1-B, 2-C, 3-A, 4-D

(A) Sodium benzoate

(B) Ethyl cellulose

(C) Eudragit

(D) Sucrose

(E) Sodium oleate

(b) 1-C, 2-D, 3-E, 4-B

(d) 1-C, 2-B, 3-D, 4-A

2.14. Match the biological listed under A to E for the following compounds:

papers are designed by the expert faculties. It may helps you in focusing your Exam preparation.

1. 1: 3-Propanediol, 2-methyl 2-propyl Carbamate

2 Chloro-10[3-(dimethylamino) propyl] Phenothiazine

5 Nitro-2-furaldeyde semicarbazone 3.

2 Methyl-5-Nitro Imidazole -ethanol

(a) 1-A, 2-B, 3-C, 4-D

(c) 1-B, 2-C, 3-A, 4-D

(A) Antimalarial

(B) Bactericidal to anaerobic and Microerophilic organisms

(C) Antibacterial

(D) Relief of anxiety and tension

(E) Tranquilizer

(b) 1-D, 2-A, 3-C, 4-B

(d) 1-E, 2-E, 3-D, 4-C



GATE - 1989

www.gdc4gpat.com

2.15. Given below are the drug A to E and the ailments for which they are recommended. Match them correctly.

- **Parkinsonism**
- 2. Hypertension
- Nasal congestion
- 4. Myasthenia gravis
- (a) 1-A, 2-B, 3-C, 4-D
- (c) 1-B, 2-C, 3-A, 4-D

- (A) Methyl dopa
- (B) Levodopa with decarboxylase inhibitor
- (C) Neostigmine
- (D) Phenyl Propanolmine
- (E) Ibuprofen
- (b) 1-B, 2-A, 3-D, 4-C
- (d) 1-C, 2-B, 3-D, 4-A

2.16. Given below are some of the drugs and their mode action in A to E. Match them correctly.

- 1. Hydralazine
- 2. Phenothiazine
- 3. Methylsergide
- Tolazmide
- (a) 1-A, 2-B, 3-C, 4-D
- (c) 1-B, 2-C, 3-A, 4-D

- (A) Vasodilator by direct action
- (B) Inhibits the Vasoconstrictor and presor effect of 5 HT
- (C) Antagonist D2 receptor of Dopamine
- (D) Stimulate the islet tissue to secrete insulin
- (E) Inhibiting the enzyme carbonic anhydrase
- (b) 1-D, 2-C, 3-B, 4-A
- (d) 1-A, 2-C, 3-B, 4-D

2.17. Given below in A to E are the names of drugs,. Appropriate tests are given below for drugs. Match them correctly.

- Alcoholic solution of α-naphthol and sulphuric acid E
- 2. Murexide test
- Para-dimethylamino Benzaldehyde 3.
- Ninhydrine 4.
- (a) 1-A, 2-B, 3-C, 4-D
- (c) 1-E, 2-C, 3-A, 4-D

- (A) Atropine
- (B) Resepine
- (C) Caffeine
- (D) Gelatin
- (E) Triticum sativum powder
- (b) 1-D, 2-C, 3-B, 4-A
- (D) 1-C, 2-B, 3-D, 4-A

2.18. Given below in A to E are the names of instruments used for the determination of the following. Match them correctly

- 1. Particle volume
- 2. Presence of Foreign particle
- Surface tension 3.
- 4. Presence of polymorph
- (a) 1-C, 2-A, 3-B, 4-E
- (c) 1-B, 2-C, 3-A, 4-D

- (A) Clarity apparatus
- (B) Du Nouy ring
- (C) Coulter counter
- (D) Compactor
- (E) Differential thermal calorimeter
- (b) 1-D, 2-C, 3-B, 4-A
- (d) 1-C, 2-B, 3-D, 4-A

Visit - www.gdconlinetest.in | Attempt 1 Free Demo Test

Email: gdcgpat037@gmail.com

YEAR PAPER

ALIAHABAD BILASPUR BHILAI RAIPUR NAGPUR HYDERABAD JAIPUR

papers are designed by the expert faculties. It may helps you in focusing your Exam preparation.



GATE - 1989

www.gdc4gpat.com

2.19. Chosse the correct starting material listed from A to E for the synthesis of the following drugs.

- 1. Cortisone
- 2. Progesterone
- 3. Testosterone
- 4. Vitamin A
- (a) 1-A, 2-B, 3-C, 4-D
- (c) 1-B, 2-C, 3-A, 4-D

- (A) Diosgenin
- (B) β-ionone
- (C) Spirostanol
- (D) Sarmentogenin
- (E) Anthracin
- (b) 1-D, 2-C, 3-B, 4-A
- (d) 1-C, 2-A, 3-A, 4-B

2.20. Given below are the types of ointment bases. Match them with the correct ointments in A to E.

- 1. Absorption base
- 2. Oleogenous base
- 3. Emulsion base
- 4. Water soluble base
- (a) 1-A, 2-B, 3-C, 4-D
- (c) 1-B, 2-E, 3-C, 4-A

- (A) Emulsifying ointmen
- (B) Hydrophilic ointment
- (C) Oily cream
- (D) Kaolin poultice
- (E) Simple ointment
- (b) 1-D, 2-A, 3-C, 4-B
- (d) 1-C, 2-B, 3-D, 4-A

DISCUSSION

PART - B

N.B.: Answer any twenty questions

If more than 20 questions are attempted only the first 20 will be considered.

Answer should not exceed 15 lines

All Question carry equal marks.

- 3. Write the structure of the following indicating to what pharmacological category they belong
 - (a) [1-dimethylamino-3-(4-Chlorophenyl 3.2-Pyridyl) Propane]
 - (b) 2. Hydroxy methylene-17 β hydroxy -17-methyl 5 α -androstan 3-one
 - (c) 2, 4-diamino-5-(3, 4, 5-trimethoxy phenyl) methyl pyridine.
- 4. (a) What is a barrier-layer cell?
 - (b) What are the different ways by which a molecule can absorb energy
- 5. Explain briefly the improved artificial method for producing Sclerotium.

papers are designed by the expert faculties. It may helps you in focusing your Exam preparation.

6. Give the characteristics of the ideal preservative for Pharmaceutical preparation.

Visit - www.gdconlinetest.in | Attempt 1 Free Demo Test

Email: gdcgpat037@gmail.com

PREVIOUS
YEAR PAPER



GATE - 1989

www.gdc4gpat.com

- 7. Give the principal of official assay of INH. Given Equations for the reactions involved.
- 8. Give the exact mode action of the following drugs:
 - (a) Dicoumarol
 - (b) Vinblastin
 - (c) Valprolic acid
- 9. Give the mode of action of the following anti arrhythmic drugs:
 - (a) Procainamide
 - (b) Propranolol
 - (c) Verapamil
- 10. Mention the various factors governing transdernal absorption of drugs.
- 11. (a) What is Hoffmann's exhaustive Methylaation?
 - (b) Show the complete step of reactions when Isoquinoline is subjected to Hoffmann's exhaustive Methylaation.
- 12. How the solid samples are prepared for the measurement of IR Absorption spectra? Why such a process is adopted
- 13. Name the various Insulin injections which are official in IP. Mention time onset and duration of action.
- 14. Give the names of the drugs, their source. And one chemical test for identification of any one important constituent in each of the following.
 - (a) Drug obtained as latex after incisions on capsule.
 - (b) Dried juice obtained from the leaves of plant belonging to Liliaceaefamily.
 - (c) A seed having action on heart
- 15. Enumerate the problems associated with use of plastic as a material for packaging Pharmaceuticals.
- 16. With the help of IR absorption readings how you can distinguish the following pairs of compounds. Predict the bands and interpret.
- **17**. Define the following:
 - (a) Liposome

(b) Polymorphism

- (c) Prodrug
- 18. Name the various methods in the preparation of micro capsules and give only the process involved in the Cocaervation phase separation technique.

Visit - www.gdconlinetest.in | Attempt 1 Free Demo Test

Email: gdcgpat037@gmail.com



www.gdc4gpat.com



- 19. Give the mode of action of following antibiotics:
 - (a) Ampicillin
 - (b) Tetracycline
 - (c) Erythromycin
- 20. What are the possible adulterants of fox glove leaves? How are they detected?
- 21. List the physic chemical factors affecting drug absorption.
- **22.** Write the equation for the following synthesis:
 - O-Chloro benzoic acid is condensed with 2, 3 Xylidine with the aid of Potassium carbonate and the resulting Potassium salt is treats with mineral acid.
- **23**. Give the possible Drug/Drug interaction of the following combination:
 - (a) Penicillin with probenecid
 - (b) Lithium carbonate with Chlorthiazide
 - (c) Levodopa with pyridoxine
- 24. (a) What concentration of Dextrose willbe used for the preparation of 100ml of Dextrose solution isotonic with blood serum. Molecular weight of Dextrose = 180
 - (b) In what proportion 80% and 30% alcohol mixed to obtain 50% alcohol.
- 25. Give the structure and specification relationship in the following compounds:
 - (a) Phenobarbital
 - (b) Amobarbital
 - (c) Cyclobarbital
 - (d) Pentobarbital
- 26. (a) An alkaloid gave E_1^1 at 310 nm = 180. The Extinction of 0.003% solution in water at 310 nm was found 0.500 (1 cm cell). Calculate the percentage of alkaloid.
 - (b) Find the HLB value of a center which has Saponificatio number 40.5 and acid number of the fatty acid 260.0
- 27. Name the endogenous neurohormones and give their structure.

End of paper

Visit - www.gdconlinetest.in | Attempt 1 Free Demo Test

Email: gdcgpat037@gmail.com




www.gdc4gpat.com



FOR GUIDANCE

ASK YOUR MORE FRIENDS TO JOIN OUR WHATSAPP CHANNEL



Visit - www.gdconlinetest.in | Attempt 1 Free Demo Test

Email: gdcgpat037@gmail.com



www.gdc4gpat.com



ANSWER KEY GATE 1989

Section - I

1 - c	2 – b	3 – c	4 – a	5 – d	6 – b	7 – b	8 – a	9 – b	10 – d
11 – b	12 – b	13 – c	14 – c	15 – b	16 – b	17 – b	18 – a	19 – b	20 – c
21 – a	22 – d	23 – c	24 – c	25 – c	26 – a	27 – d	28 – c	29 – b	30 – b
								39 – b	

Section - II

2.1 - a	2.2 - b	2.3 - d	2.4 - a	2.5 - c	2.6 - b	2.7 - a	2.8 - b	2.9 - d
2.10 - a	2.11 - c	2.12 - c	2.13 - b	2.14 - d	2.15 - b	2.16 - d	2.17 - c	2.18 – a
2.19 - d	2.20 – c							



STAY CONNECTED WITH US AS LOT OF THINGS ARE COMING FOR YOU

ALL THE BEST FOR YOUR SMART PREPARATION
BEST REGARDS FROM
TEAM GDC